

FITC-Labeled Human CD37 Protein, Fc Tag

Catalog # CD7-HF26x



BIOSYSTEMS
Acro

Surprise Inside!

Synonym

CD37, Tspan-26, TSPAN26

Source

FITC-Labeled Human CD37, Fc Tag (CD7-HF26x) is expressed from human 293 cells (HEK293). It contains AA Arg 112 - Asn 241 (Accession # [P11049-1](#)). It is the FITC labeled form of Human CD37, Fc Tag (Cat. No. CD7-H525a).

Predicted N-terminus: Arg 112

Molecular Characterization

CD37(Arg 112 - Asn 241) P11049-1	Fc(Pro 100 - Lys 330) P01857
-------------------------------------	---------------------------------

[Other Tags and Version Biotin & Other Labeled Version](#)

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 41.3 kDa. The protein migrates as 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Conjugate

FITC

Excitation source: 488 nm spectral line, argon-ion laser

Excitation Wavelength: 488 nm

Emission Wavelength: 535 nm

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with FITC using standard chemical labeling method. The residual FITC is removed by molecular sieve treatment during purification process.

Protein Ratio

The FITC to protein molar ratio is **1.5-3.5**.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE



FITC-Labeled Human CD37, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Background

Leukocyte antigen CD37 (CD37) is also known as Tetraspanin-26 (Tspan-26) and TSPAN26, which is a member of the transmembrane 4 superfamily or tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins encoded by CD37 gene mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. CD37 is a cell surface glycoprotein that is known to complex with integrins and other transmembrane 4 superfamily proteins. CD37 may play a role in T-cell-B-cell interactions.

Discounts, Gifts,
and more!



www.acrobiosystems.com

11/20/2025