

**Synonym**

Basigin,BSG,5F7,CD147,EMMPRIIN,M6,OK,TCSF

Source

Biotinylated Human EMMPRIN, His,Avitag(CD7-M82E8) is expressed from human 293 cells (HEK293). It contains AA Ala 22 - Ala 211 (Accession # [NP_001070652.1](#)).

Predicted N-terminus: Ala 22

Molecular Characterization

EMMPRIIN(Ala 22 - Ala 211) NP_001070652.1	Poly-his	Avi
--	----------	-----

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 24.3 kDa. The protein migrates as 33-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.


Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE

Biotinylated Human EMMPRIN, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

SEC-MALS

The purity of Biotinylated Human EMMPRIN, His,Avitag (Cat. No. CD7-M82E8) is more than 85% and the molecular weight of this protein is around 28-42 kDa verified by SEC-MALS.

[Report](#)**Background****Discounts, Gifts,
and more!**

Biotinylated Mouse EMMPRIN / CD147 Protein, His,Avitag™ (MALS verified)

Catalog # CD7-M82E8



BIOSYSTEMS
Acro

CD147, also known as Basigin (BSG), or extracellular matrix metalloproteinase inducer (EMMPRIN). The human basigin protein contains 269 amino acids that form two heavily glycosylated C2 type immunoglobulin-like domains at the N-terminal extracellular portion. A second form of basigin has also been characterized that contains one additional immunoglobulin-like domain in its extracellular portion. As members of the immunoglobulin superfamily play fundamental roles in intercellular recognition involved in various immunologic phenomena, differentiation, and development, basigin is thought also to play a role in intercellular recognition and regulate several distinct functions, such as spermatogenesis, expression of the monocarboxylate transporter and the responsiveness of lymphocytes. Basigin is a type I integral membrane receptor that has many ligands, including the cyclophilin (CyP) proteins Cyp-A and CyP-B and certain integrins.

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

8/11/2025