

APC-Labeled Monoclonal Anti-Human CD28 Antibody, Mouse IgG1 (0.03% Proclin)

Catalog # CD8-AHFC1



BIOSYSTEMS
Acro

Source

Monoclonal Anti-Human CD28 Antibody, Mouse IgG1 is a monoclonal antibody recombinantly expressed from human 293 cells (HEK293), which provides higher batch consistency and long term security of supply.

Application

Flow Cytometry (Evaluation of the expression of CD28 on Human cells).

Species

Mouse

Isotype

Mouse IgG1 | Mouse Kappa

Specificity

This product is a specific antibody specifically reacts with CD28 protein.

Reactivity

Human

Immunogen

Purified Human CD28 Protein

Conjugate

APC

Excitation Wavelength: 640 nm

Emission Wavelength: 661 nm

Recommended Dilution

1:50

Formulation

Lyophilized from a 0.22 µm-filtered solution in PBS (pH 7.4) containing 0.03% ProClin 300 and 0.2% BSA, with trehalose as protectant.

Please contact us for customized product forms or formulations.

Reconstitution

Please refer to the Certificate of Analysis (CoA) for specific instructions.

For best performance, we strongly recommend following the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored in a lyophilized state at -20°C or lower.

Please protect from light and avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 24 months in lyophilized state;
- -70°C for 12 months after reconstitution;
- 2-8 °C for 12 months after reconstitution.


ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

Bioactivity-FACS



Flow cytometric analysis of Human peripheral blood lymphocytes respectively staining with APC-Labeled Monoclonal Anti-Human CD28 Antibody, Mouse IgG1 (Cat. No. CD8-AHFC1) at 1:50 dilution (2 μ L of the antibody stock solution corresponds to labeling of 1e6 cells in a final volume of 100 μ L), compared with isotype control antibody. APC signal was used to evaluate the binding activity (QC tested).



Non-specificity of APC-Labeled Monoclonal Anti-Human CD28 Antibody, Mouse IgG1 (Cat. No. CD8-AHFC1) binding to CD3- cells present in human PBMC. Human PBMCs were simultaneously stained with FITC anti-human CD3 Antibody and APC-Labeled Monoclonal Anti-Human CD28 Antibody, Mouse IgG1 (2 μ L of the antibody stock solution corresponds to labeling of 5e5 cells in a final volume of 100 μ L), washed and then analyzed with FACS. FITC negative signals and APC positive signals were used to evaluate the non-specific binding activity to human CD3- cells (Routinely tested).

Background

T-cell-specific surface glycoprotein CD28 is also known as TP44, is a single-pass type I membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. is one of the molecules expressed on T cells that provide co-stimulatory signals, which are required for T cell activation. CD28 is the receptor for CD80 (B7.1) and CD86 (B7.2). When activated by Toll-like receptor ligands, the CD80 expression is upregulated in antigen presenting cells (APCs). The CD86 expression on antigen presenting cells is constitutive. CD28 is the only B7 receptor constitutively expressed on naive T cells.



Discounts, Gifts,
and more!

www.acrobiosystems.com

