



## Background

Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4) is a CK20 primary antibody for IHC application. CK20 belongs to cytokeratin family and typically expressed in the epithelial cells of the gastrointestinal and urinary tracts. CK20 serves as a critical tumor biomarker in the clinical practice in a variety of tumors in the colon and bladder.

## Key parameter

Host Species	Rabbit
Isotype	IgG
Clone	5D4
Application	IHC
Property	1: 400
State	Liquid
Clonality	Monoclonal
Research Field	Cancer Drug Targets
Source	Rabbit
Synonym	Cytokeratin 20

## Experiment Protocol


LEICA Experiment Protocol(Manual)

## Storage


Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. The shelf life is 30 days from the date of opening.

## Typical Data

### Control Sample



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4) Human Colorectal Cancer, 4X



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4) Human Colorectal Cancer, 20X

**Immunohistochemical analysis of paraffin embedded human colorectal cancer labelled with HGS-S602 at 1/400 dilution.**

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Cancer Sample

Discounts, Gifts,  
and more!




# Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4)


Catalog # HGS-S602



BIOSYSTEMS  
**Acro**



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4) Human Urothelial Carcinoma, 4X



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-CK20 Antibody, Rabbit (5D4) Human Urothelial Carcinoma, 20X

## Immunohistochemical analysis of paraffin embedded human urothelial carcinoma labelled with HGS-S602 at 1/400 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Discounts, Gifts,  
and more!



➤ [www.acrobiosystems.com](http://www.acrobiosystems.com)

6/25/2025