



## Background

Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5) is a TTF-1 primary antibody for IHC application. TTF-1 is expressed in the epithelial cells and positively expressed different types of cancers. TTF-1 serves as a critical tumor biomarker in the clinical practice in a variety of tumors in the lung.

## Key parameter

Host Species	Rabbit
Isotype	IgG
Clone	1A5
Application	IHC
Property	1: 400
State	Liquid
Positive Control	Human Lung Tissue
Clonality	Monoclonal
Research Field	Cancer Drug Targets
Source	Rabbit
Synonym	TTF-1

## Experiment Protocol


LEICA Experiment Protocol(Manual)

## Storage


Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. The shelf life is 30 days from the date of opening.

## Typical Data

### Control Sample



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5) Human Lung Adenocarcinoma, 4X



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5) Human Lung Adenocarcinoma, 20X

**Immunohistochemical analysis of paraffin embedded human lung adenocarcinoma labelled with HGS-S606 at 1/400 dilution.**

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Cancer Sample

Discounts, Gifts,  
and more!




# Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5)


Catalog # HGS-S606



BIOSYSTEMS  
**Acro**



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5) Human Thyroid Carcinoma, 4X



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Recombinant Monoclonal Anti-TTF-1 Antibody, Rabbit (1A5) Human Thyroid Carcinoma, 20X

**Immunohistochemical analysis of paraffin embedded human thyroid carcinoma labelled with HGS-S606 at 1/400 dilution.**

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Discounts, Gifts,  
and more!



➤ [www.acrobiosystems.com](http://www.acrobiosystems.com)