

Human IL-7 Protein, premium grade

Catalog # IL7-H4219



BIOSYSTEMS
Acro

Features and Advantages

- Seamless Transition from Preclinical to Clinical: Same cell clone, process, and QC as GMP products.
- Enhanced Production Standards: AOF raw materials, pharma-grade excipients, Grade B+A (ISO 5) environment.
- Comprehensive Quality Control: Additional strict testing for process impurities, sterility, and mycoplasma.
- Cost-Effective Solution: GMP-comparable quality at an economical price for early development.
- Risk Mitigation: Enhanced safety through rigorous impurity control and comprehensive testing.

Synonyms

IL7, Interleukin-7

Source

Human IL-7, premium grade (IL7-H4219) is expressed from human 293 cells (HEK293). It contains AA Asp 26 - His 177 (Accession # [P13232-1](#)).

Predicted N-terminus: Asp 26

It is produced under our rigorous quality control system that incorporates a comprehensive set of tests including sterility and endotoxin tests. Product performance is carefully validated and tested for compatibility for cell culture use or any other applications in the early preclinical stage.

GMP-L07H24 is the GMP version of this IL7-H4219. These two proteins display indistinguishable performance profiles, thereby ensuring a seamless transition for end users from early preclinical stage to later clinical phases.

Molecular Characterization

IL-7(Asp 26 - His 177)
P13232-1

This protein carries no "tag".

The protein has a calculated MW of 17.4 kDa. The protein migrates as 23 kDa and 28 kDa (± 3 kDa) when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per μg by the LAL method / rFC method.

Host Cell Protein

<0.5 ng/ μg of protein tested by ELISA.

Host Cell DNA

<0.02 ng/ μg of protein tested by DNA Fluorescent Staining method.

Sterility

Negative

Mycoplasma

Negative

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 μm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

Quality Description

ACRO's Research-grade products are suitable for a wide range of cell culture applications, particularly for research use in academic institutions. These products are sterilized by filtration, followed by lyophilization where applicable. Typical specifications include endotoxin levels of <0.01 EU/μg and purities >95%. Biological activity is calibrated against WHO/NIBSC standards when available.


ACRO's Premium-grade (Pre-GMP) products are characterized by their high quality and enhanced safety profiles, making them ideal for early-stage discovery and manufacturing processes in cell therapy companies. A key advantage is their seamless transition to corresponding GMP-grade versions. Biological activity is calibrated against WHO/NIBSC standards when available. Typical specifications include endotoxin levels of <0.01 EU/μg and purities >95%. In addition, rigorous testing is conducted to ensure the absence of mycoplasma, HCD, and HCP, thereby guaranteeing product safety.

SDS-PAGE




Human IL-7, premium grade on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-CELL BASE



Human IL-7, premium grade (Cat. No. IL7-H4219) stimulates proliferation of PHA-P-activated human peripheral blood mononuclear cell (PBMC). The specific activity of Human IL-7, premium grade is > 1.00×10⁸ IU/mg, which is calibrated against human IL-7 WHO International Standard (NIBSC code: 90/530) (QC tested).

Bioactivity-Stability



Cell-based assay demonstrates batch-to-batch consistency between Acro's GMP and PG IL-7.

Background

Interleukin 7 is also known as IL7, IL-7, and is a hematopoietic growth factor secreted by stromal cells in the red marrow and thymus. It is also produced by keratinocytes, dendritic cells, hepatocytes, neurons, and epithelial cells, but is not produced by lymphocytes. IL-7 stimulates the differentiation of multipotent (pluripotent) hematopoietic stem cells into lymphoid progenitor cells, It also stimulates proliferation of all cells in the lymphoid lineage (B cells, T cells and NK cells). It is important for proliferation during certain stages of B-cell maturation, T and NK cell survival, development and homeostasis. IL-7 is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. IL-7 binds to the IL-7 receptor, a heterodimer consisting of Interleukin-7 receptor alpha and common gamma chain receptor. Il-7 promotes hematological malignancies (acute lymphoblastic leukemia, T cell lymphoma). Elevated levels of IL-7 have also been detected in the plasma of HIV-infected patients. IL-7 as an immunotherapy agent has been examined in many pre-clinical animal studies and more recently in human clinical trials for various malignancies and during HIV infection. IL-7 could also be beneficial in improving immune recovery after allogenic stem cell transplant.



www.acrobiosystems.com