

Synonym

IL-12 R beta 1, IL12RB1, CD212, IL12R, IL12RB, IL-12RB1

Source

Cynomolgus IL-12 R beta 1 Protein, Fc Tag (ILB-C5253) is expressed from human 293 cells (HEK293). It contains AA Cys 24 - Asp 545 (Accession # [G7PWY9-1](#)).  
Predicted N-terminus: Cys 24

Molecular Characterization

|                                        |                                 |
|----------------------------------------|---------------------------------|
| IL-12RB1(Cys 24 - Asp 545)<br>G7PWY9-1 | Fc(Pro 100 - Lys 330)<br>P01857 |
|----------------------------------------|---------------------------------|

[Other Tags and Version Biotin & Other Labeled Version](#)

This protein carries a human IgG1 Fc tag at the C-terminus.  
The protein has a calculated MW of 83.9 kDa. The protein migrates as 110-120 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.  
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.  
**For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.**

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.  
**Please avoid repeated freeze-thaw cycles.**  
This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)


SDS-PAGE



Cynomolgus IL-12 R beta 1 Protein, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein

is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

## Bioactivity-ELISA



Immobilized Human IL-12B&IL-12A Heterodimer Protein, His Tag&Flag Tag (Cat. No. IL2-H4210) at 5 µg/mL (100 µL/well) can bind Cynomolgus IL-12 R beta 1 Protein, Fc Tag (Cat. No. ILB-C5253) with a linear range of 0.005-0.156 µg/mL (QC tested).

## Background

The human IL-12 R subunit is a member of the cytokine receptor superfamily and the functional high-affinity IL-12R is composed of at least two beta-type cytokine receptor subunits, each independently exhibiting a low affinity for IL-12. IL-12 R beta 1 (Interleukin-12 receptor subunit beta-1) is also known as IL-12RB1, CD212. Functions as an interleukin receptor which binds interleukin-12 with low affinity and is involved in IL12 transduction. Associated with IL12RB2 it forms a functional, high affinity receptor for IL12. Associates also with IL23R to form the interleukin-23 receptor which functions in IL23 signal transduction probably through activation of the Jak-Stat signaling cascade.

Discounts, Gifts,  
and more!



Acro



[www.acrobiosystems.com](http://www.acrobiosystems.com)