

Human Insulin R / CD220 Protein, His Tag (MALs & SPR verified)

Catalog # INR-H52H3



Synonyms

INSR, Insulin receptor, IR, CD220

Source

Human Insulin R Protein, His Tag (INR-H52H3) is expressed from human 293 cells (HEK293). It contains AA His 28 - Lys 956 (Accession # [P06213-1](#)).

Molecular Characterization

Insulin R(His 28 - Lys 956) P06213-1	Poly-his
---	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 107.8 kDa. The protein migrates as 35-45 kDa and 95-180 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

The protein is designed as a dimer.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE




Human Insulin R Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS



The purity of Human Insulin R Protein, His Tag (Cat. No. INR-H52H3) is more than 90% and the molecular weight of this protein is around 250-300 kDa verified by SEC-MALS.

Bioactivity-SPR



Human Insulin R Protein, His Tag (Cat. No. INR-H52H3) immobilized on CM5 Chip can bind Human IGF-II, Fc Tag (Cat. No. IG2-H4260) with an affinity constant of 1.81 μ M as determined in a SPR assay (Biacore 8K) (QC tested).

Background

Insulin receptor (INSR) is also known as CD antigen CD220, which can be cleaved into the following 2 chains: Insulin receptor subunit alpha and Insulin receptor subunit beta. INSR is a tetramer of 2 alpha and 2 beta chains linked by disulfide bonds. The alpha chains carry the insulin-binding regions, while the beta chains carry the kinase domain. Forms a hybrid receptor with IGF1R, the hybrid is a tetramer consisting of 1 alpha chain and 1 beta chain of INSR and 1 alpha chain and 1 beta chain of IGF1R. In addition to binding insulin, the insulin receptor can bind insulin-like growth factors (IGFI and IGFII). Isoform Short of INSR has a higher affinity for IGFII binding. When present in a hybrid receptor with IGF1R, INSR binds IGF1.



www.acrobiosystems.com