



Synonym

betaKlotho, beta-klotho, BKL, KLB, klotho beta like, klotho beta-like protein

Source

Biotinylated Human Klotho beta Protein, His Tag, primary amine labeling (KLB-H82H3) is expressed from human 293 cells (HEK293). It contains AA Met 30 - Leu 997 (Accession # [Q86Z14-1](#)).

Predicted N-terminus: Met 30

Molecular Characterization

KLB(Met 30 - Leu 997)
Q86Z14-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 112.9 kDa. The protein migrates as 130-150 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

The primary amines in the side chains of lysine residues and the N-terminus of the protein are conjugated with biotins using standard chemical labeling method. A standard biotin reagent (13.5 angstroms) is used in this product.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE



Biotinylated Human Klotho beta Protein, His Tag, primary amine labeling on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA



Immobilized Human FGF19, Fc Tag (Cat. No. FG9-H5253) at 5 µg/mL (100 µL/well) can bind Biotinylated Human Klotho beta Protein, His Tag, primary amine labeling (Cat. No. KLB-H82H3) with a linear range of 0.02-0.625 µg/mL (QC tested).

Background

KLB (Klotho Beta) is a Protein Coding gene. Among its related pathways are RET signaling and HIV Life Cycle. GO annotations related to this gene include hydrolase activity, hydrolyzing O-glycosyl compounds and fibroblast growth factor binding. An important paralog of this gene is KL. Klotho Beta is a regulator in multiple metabolic processes, while its role in cancer remains unclear. We found the expression of β Klotho was down-regulated in human hepatocellular carcinoma tissues compared with that in paired adjacent non-tumourous liver tissues. Hepatoma cells also showed decreased expression of β Klotho compared with normal hepatocyte cells. Reintroduction of β Klotho into hepatoma cells inhibited their proliferation.

Discounts, Gifts,
and more!



www.acrobiosystems.com

