

Mouse HGF R / c-MET Protein, Fc Tag (MALS verified)

Catalog # MET-M5254



Synonym

MET, AUTS9, HGFR, RCCP2, c-Met

Source

Mouse HGF R, Fc Tag (MET-M5254) is expressed from human 293 cells (HEK293). It contains AA Glu 25 - Ala 931 (Accession # [P16056-1](#)).
Predicted N-terminus: Glu 25

Molecular Characterization

HGF R(Glu 25 - Ala 931) P16056-1	Fc(Pro 100 - Lys 330) P01857
-------------------------------------	---------------------------------

Other Tags and Version [Biotin & Other Labeled Version](#)

Mouse HGF R, Fc Tag is fused with Fc fragment of human IgG1 at the C-terminus. The mature form of HGF R is a disulfide-linked heterodimer composed of proteolytically cleaved α and β chain. Each α and β chain has a calculated MW of 32.1 kDa (α chain) and 95.1 kDa (β chain Fc chimera). The protein migrates as 40-45 kDa (α chain) and 125-135 kDa (β chain Fc chimera) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per μ g by the LAL method / rFC method.

Purity

- >95% as determined by SDS-PAGE.
- >95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:

- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.


ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE




SEC-MALS



Mouse HGF R, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

The purity of Mouse HGF R, Fc Tag (Cat. No. MET-M5254) is more than 95% and the molecular weight of this protein is around 254-311 kDa verified by SEC-MALS.

Bioactivity-ELISA



Immobilized Human HGF, His Tag (Cat. No. HGF-H52H3) at 2 µg/mL (100 µL/well) can bind Mouse HGF R, Fc Tag (Cat. No. MET-M5254) with a linear range of 0.2-10 ng/mL (QC tested).

Background

Hepatocyte growth factor receptor (HGFR) is also known as mesenchymal-epithelial transition factor (MET), c-Met, and is a glycosylated receptor tyrosine kinase that plays a central role in epithelial morphogenesis and cancer development. HGFR protein possesses tyrosine-kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. HGFR is normally expressed by cells of epithelial origin, while expression of HGF is restricted to cells of mesenchymal origin. Upon HGF stimulation, HGFR induces several biological responses that collectively give rise to a program known as invasive growth. Abnormal HGFR activation in cancer correlates with poor prognosis, where aberrantly active HGFR triggers tumor growth, formation of new blood vessels (angiogenesis) that supply the tumor with nutrients, and cancer spread to other organs (metastasis). HGFR is deregulated in many types of human malignancies, including cancers of kidney, liver, stomach, breast, and brain. Normally, only stem cells and progenitor cells express HGFR, However, cancer stem cells are thought to hijack the ability of normal stem cells to express HGFR, and thus become the cause of cancer persistence and spread to other sites in the body. Various mutations in the HGFR gene are associated with papillary renal carcinoma. HGFR mediates a complex program known as invasive growth. Activation of HGFR triggers mitogenesis, and morphogenesis.

Discounts, Gifts,
and more!





www.acrobiosystems.com