

Synonym

Mucin 1,MUC1,CD227,EMA,H23AG,KL-6,MAM6,MUC-1,SEC,MUC-1,X,MUC1,ZD,PEM,PEMT,PUM,CA15-3,Episialin

Source

Human Mucin-1 (890-1158) Protein, Fc Tag(MU1-H5258) is expressed from human 293 cells (HEK293). It contains AA Ser 890 - Gly 1158 (Accession # [P15941-1](#)).
Predicted N-terminus: Ser 890

Molecular Characterization

Mucin-1(Ser 890 - Gly 1158) P15941-1	Fc(Pro 100 - Lys 330) P01857
---	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.
The protein has a calculated MW of 54.3 kDa. The protein migrates as 40-43 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE).

Endotoxin

Less than 0.1 EU per µg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.
>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH7.5 with trehalose as protectant.
Contact us for customized product form or formulation.

Reconstitution


Please see Certificate of Analysis for specific instructions.
For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.
Please avoid repeated freeze-thaw cycles.
This product is stable after storage at:


- 20°C to -70°C for 12 months in lyophilized state;
- 70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



Human Mucin-1 (890-1158) Protein, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).


SEC-MALS



The purity of Human Mucin-1 (890-1158) Protein, Fc Tag (Cat. No. MU1-H5258) is more than 90% and the molecular weight of this protein is around 160-185 kDa verified by SEC-MALS.
[Report](#)

Bioactivity-ELISA





Immobilized Human Mucin-1 (890-1158) Protein, Fc Tag (Cat. No. MU1-H5258) at 1 µg/mL (100 µL/well) can bind Anti-Mucin-1 Antibody, Mouse IgG1 with a linear range of 0.1-4 ng/mL (QC tested).

Background

This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants

