

Rat Nectin-4 Protein, His Tag (MALS verified)

Catalog # NE4-R52H3



Synonyms

NECTIN4, LNIR, PRR4, PVRL4

Source

Rat Nectin-4 Protein, His Tag (NE4-R52H3) is expressed from human 293 cells (HEK293). It contains AA Tyr 28 - Val 348 (Accession # [NP_001102546.1](#)).

Predicted N-terminus: Tyr 28

Molecular Characterization

Nectin-4(Tyr 28 - Val 348) NP 001102546.1	Poly-his
--	----------

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 36.6 kDa. The protein migrates as 35-40 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Purity

>90% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE




Rat Nectin-4 Protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

SEC-MALS



The purity of Rat Nectin-4 Protein, His Tag (Cat. No. NE4-R52H3) is more than 95% and the molecular weight of this protein is around 36-50 kDa verified by SEC-MALS.

Bioactivity-ELISA



Immobilized Rat Nectin-4 Protein, His Tag (Cat. No. NE4-R52H3) at 1 µg/mL (100 µL/well) can bind Mouse Anti-Nectin-4 Antibody, Mouse IgG1 with a linear range of 0.01-2.5 µg/mL (QC tested).

Background

Nectin-4 (gene name PVRL4, poliovirus receptor-like 4) is a 66 kDa type I transmembrane glycoprotein belonging to the Nectin family of Ig superfamily proteins. Nectins are cell adhesion molecules that play a key role in various biological processes such as polarity, proliferation, differentiation and migration, for epithelial, endothelial, immune and neuronal cells, during development and adult life. Nectin-4 is a tumor-associated antigen in 50%, 49% and 86% of breast, ovarian and lung carcinomas, respectively, mostly on tumors of bad prognosis. Its expression is not detected in the corresponding normal tissues.

Discounts, Gifts,
and more!



www.acrobiosystems.com