

Human IL-12B / NKSF2 / p40 Protein, His Tag

Catalog # NK2-H52H7



BIOSYSTEMS
Acro

Synonyms

IL12B, CLMF, CLMF2, NKSF, NKSF2, p40

Source

Human IL-12B, His Tag (NK2-H52H7) is expressed from human 293 cells (HEK293). It contains AA Ile 23 - Ser 328 (Accession # [P29460-1](#)).

Predicted N-terminus: Ile 23

Molecular Characterization

IL-12B(Ile 23 - Ser 328)
P29460-1

Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 36.6 kDa. The protein migrates as 37-45 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE




Human IL-12B, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA




Immobilized Monoclonal Anti-Human IL23A&IL12B P40 domain Antibody, Human IgG1 at 2 µg/mL (100 µL/well) can bind Human IL-12B, His Tag (Cat. No. NK2-H52H7) with a linear range of 1-8 ng/mL (QC tested).

Bioactivity-BLI



Loaded Human IL-12 R beta 1, Fc Tag (Cat. No. ILB-H5255) on Protein A Biosensor, can bind Human IL-12B, His Tag (Cat. No. NK2-H52H7) with an affinity constant of 3.78 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded Anti-IL23A & IL12B MAb (P40 domain) on Protein A Biosensor, can bind Human IL-12B, His Tag (Cat. No. NK2-H52H7) with an affinity constant of 0.279 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

Interleukin-12 (IL12) is also known as natural killer cell stimulatory factor (NKSF), cytotoxic lymphocyte maturation factor (CLMF) , is a heterodimeric cytokine encoded by two separate genes, IL-12A (p35) and IL-12B (p40). IL12 is naturally produced by dendritic cells, macrophages and human B-lymphoblastoid cells (NC-37) in response to antigenic stimulation. IL-12 is involved in the differentiation of naive T cells into Th0 cells and plays an important role in the activities of natural

killer cells and T lymphocytes.IL-12 also has anti-angiogenic activity, which means it can block the formation of new blood vessels. Interleukin-12 subunit beta (IL12B) also known as NKSF2, CLMF2 and P40. Interleukin-12 subunit beta has been shown to interact with IL23. A large excess of monomeric IL12B is also secreted by the cells producing IL12, and exhibits no demonstrable biological activity. Overexpression of IL12B gene has been shown to be associated with the pathogenesis of multiple sclerosis. In addition, studies have revealed that the promoter polymorphism of this gene is implicated in the severity of atopic and non-atopic asthma in children.



www.acrobiosystems.com