

Synonyms

NKG2D, CD314, KLRK1, NK cell receptor D

Source

Human NKG2D, Mouse IgG2a Fc Tag (NKD-H5259) is expressed from human 293 cells (HEK293). It contains AA Ile 73 - Val 216 (Accession # [P26718-1](#)).

Predicted N-terminus: Glu

Molecular Characterization

|                                 |                                     |
|---------------------------------|-------------------------------------|
| mFc(Glu 98 - Lys 330)<br>P01863 | NKG2D(Ile 73 - Val 216)<br>P26718-1 |
|---------------------------------|-------------------------------------|

This protein carries a mouse IgG2a Fc tag at the N-terminus.

The protein has a calculated MW of 43.5 kDa. The protein migrates as 50-60 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.1 EU per µg by the LAL method / rFC method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE



Human NKG2D, Mouse IgG2a Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA



Immobilized Human MICA, His Tag (Cat. No. MIA-H5221) at 1 µg/mL (100 µL/well) can bind Human NKG2D, Mouse IgG2a Fc Tag (Cat. No. NKD-H5259) with a linear range of 0.02-0.313 µg/mL (QC tested).

Background

NKG2D is a transmembrane protein belonging to the CD94/NKG2 family of C-type lectin-like receptors, also known as KLRK1, CD314, D12S2489E, KLR and killer cell lectin like receptor K1. NKG2D itself forms a homodimer whose ectodomains serve for ligand binding. NKG2D is a major recognition receptor for the detection and elimination of transformed and infected cells as its ligands are induced during cellular stress, either as a result of infection or genomic stress such as in cancer. In NK cells, NKG2D serves as an activating receptor, which itself is able to trigger cytotoxicity. The function of NKG2D on CD8+ T cells is to send co-stimulatory signals to activate them.



[www.acrobiosystems.com](http://www.acrobiosystems.com)