

Human PAR1 Full Length Protein, Flag,His Tag (Detergent)

Catalog # PA1-H52D3



BIOSYSTEMS
Acro

Application

- Immunization
- Affinity testing (SPR / BLI)
- Screening ([ELISA](#))

Synonym


F2R, coagulation factor II thrombin receptor, CF2R, HTR, PAR-1, PAR1, TR

Source

Detergent Human PAR1 Full Length Protein, Flag,His Tag (PA1-H52D3) is expressed from human 293 cells (HEK293). It contains AA Ser 42 - Ser 395 (Accession # [P25116-1](#)).

Predicted N-terminus: Asp


Molecular Characterization



This protein carries a flag tag at the N-terminus and a polyhistidine tag at the C-terminus.


The protein has a calculated MW of 58.7 kDa. The protein migrates as 55-180 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

• Structure



Each full-length transmembrane protein is encapsulated by detergent micelle.

• Synthesis Process



Through **mild detergent micelle extraction**, dedicated solubilization, and stringent affinity purification, we maximally retain the native transmembrane conformation of full-length transmembrane proteins throughout downstream purification. This workflow yields purified, detergent-solubilized proteins free of unrelated membrane impurities and exogenous scaffold proteins, with **exceptional purity, accurate quantitation**, and well-preserved biological activity—ideal for immunization, binding assays in antibody discovery and screening (ELISA) and affinity testing (SPR and BLI), etc.

Endotoxin

Less than 1.0 EU per μg by the LAL method / rFC method.

Purity

>90% as determined by SDS-PAGE.

Formulation

This product is not suitable for cell based experiments due to cytotoxicity of DDM.

DDM and CHS are INDISPENSABLE to keep membrane protein soluble and active, under no circumstances should you remove DDM and

CHS.

DDM/CHS buffer (DC-11) is sold separately and not included in protein, and please contact us if you need the buffer.

If glycerol is not compatible with your application, remove glycerol just before the immediate experiment, and NEVER store glycerol-free protein solution.

Supplied as 0.2 µm filtered solution in 50 mM HEPES, 150 mM NaCl, DDM, CHS, pH7.5 with glycerol as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.

Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:


- The product **MUST** be stored at -70°C or lower upon receipt;
- -70°C for 12 months under sterile conditions.

ACRO Quality Management System

- [QMS\(ISO, GMP\)](#)
- [Quality Advantages](#)
- [Quality Control Process](#)


*The DDM/CHS buffer (Cat. No. [DC-11](#)) is sold separately and not included in protein, you can follow [the link](#) for product information.

SDS-PAGE



Human PAR1 Full Length Protein, Flag,His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA



Immobilized Human PAR1 Full Length Protein, Flag,His Tag (Cat. No. PA1-H52D3) at 1 µg/mL (100 µL/well) can bind PAR1 Monoclonal Antibody with a linear range of 0.2-5 ng/mL (QC tested).

Background

PAR1, or Protease-Activated Receptor 1, is a key protein on cell surfaces that responds to thrombin, an enzyme involved in blood clotting. When activated, PAR1 triggers critical cellular responses, including platelet aggregation and inflammation. It plays a vital role in hemostasis and thrombosis, making it an important target for antiplatelet therapies.



www.acrobiosystems.com

