


**Synonym**

PD-L1,CD274,B7-H1,PDCD1L1,PDCD1LG1

**Source**

Biotinylated Mouse PD-L1 Protein, His,Avitag(PDL-M82E7) is expressed from human 293 cells (HEK293). It contains AA Phe 19 - Thr 238 (Accession # [Q9EP73](#)).

Predicted N-terminus: Phe 19

**Molecular Characterization**

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 28.3 kDa. The protein migrates as 39-54 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

**Labeling**

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*

**Protein Ratio**

Passed as determined by the HABA assay / binding ELISA.

**Purity**

&gt;90% as determined by SDS-PAGE.

&gt;90% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*


**Storage**

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.



**SDS-PAGE**

Biotinylated Mouse PD-L1 Protein, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

**SEC-MALS**

The purity of Biotinylated Mouse PD-L1 Protein, His,Avitag (Cat. No. PDL-M82E7) is more than 90% and the molecular weight of this protein is around 35-50 kDa verified by SEC-MALS.

[Report](#)**Bioactivity-ELISA****Discounts, Gifts,  
and more!**



Immobilized Anti-PD-L1 Neutralizing Antibody at 2 µg/mL (100 µL/well) can bind Biotinylated Mouse PD-L1 Protein, His,Avitag (Cat. No. PDL-M82E7) with a linear range of 0.1-2 ng/mL (QC tested).

Immobilized Biotinylated Mouse PD-L1 Protein, His,Avitag (Cat. No. PDL-M82E7) at 1 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Mouse PD-1, Fc Tag (Cat. No. PD1-M5259) with a linear range of 0.3-39 ng/mL (Routinely tested).

## Background

Programmed cell death 1 ligand 1 (PDL1) is also known as B7-H, B7H1, MGC142294, MGC142296, PD-L1, PDCD1L1 and PDCD1LG1, which is a member of the growing B7 family of immune molecules and is involved in the regulation of cellular and humoral immune responses. PDL1 is a cell surface immunoglobulin superfamily with two Ig-like domains within the extracellular region and a short cytoplasmic domain. This protein is broadly expressed in the majority of peripheral tissues as well as hematopoietic cells. Interaction between PDL1 and its receptors belonging to the CD28 family of molecules provide both stimulatory and inhibitory signals in regulating T cell activation and tolerance. PDL1 may inhibit ongoing T-cell responses by inducing apoptosis and arresting cell-cycle progression.

Discounts, Gifts,  
and more!

