

Biotinylated Rat TNFSF11 / RANKL / CD254 Protein, His,Avitag™

Catalog # RAL-R82Q9



Synonyms


RANKL, CD254, TRANCE, OPGL, ODF

Source

Biotinylated Rat TNFSF11, His,Avitag (RAL-R82Q9) is expressed from human 293 cells (HEK293). It contains AA Leu 69 - Asp 318 (Accession # [Q9ESE2-1](#)).

Predicted N-terminus: His

Molecular Characterization



This protein carries a polyhistidine tag at the N-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 31.8 kDa. The protein migrates as 35-37 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Purity

>90% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Shipping and Storage

This product is shipped at ambient temperature.

For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

ACRO Quality Management System


- [QMS\(ISO, GMP\)](#).
- [Quality Advantages](#)
- [Quality Control Process](#)

SDS-PAGE




Biotinylated Rat TNFSF11, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

Bioactivity-ELISA



Immobilized Human RANK, Mouse IgG2a Fc Tag, low endotoxin (Cat. No. RAK-H5251) at 1 µg/mL (100 µL/well) can bind Biotinylated Rat TNFSF11, His,Avitag (Cat. No. RAL-R82Q9) with a linear range of 0.5-16 ng/mL (QC tested).



Immobilized Biotinylated Rat TNFSF11, His,Avitag (Cat. No. RAL-R82Q9) at 2 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (1 µg/well) plate can bind Human Osteoprotegerin, Fc Tag (Cat. No. TNB-H5259) with a linear range of 0.1-16 ng/mL (Routinely tested).

Background

Receptor activator of nuclear factor kappa-B ligand (RANKL), also known as tumor necrosis factor ligand superfamily member 11 (TNFSF11), TNF-related activation-induced cytokine (TRANCE), osteoprotegerin ligand (OPGL), and osteoclast differentiation factor (ODF), is known as a type II membrane protein and is a member of the tumor necrosis factor (TNF) superfamily. RANKL, through its ability to stimulate osteoclast formation and activity, is a critical mediator of bone resorption and overall bone density. Some findings also suggestion some cancer cells, particularly prostate cancer cells, can activate an increase in bone remodeling and ultimately increase overall bone production.[17] This increase in bone remodeling and bone production increases the overall growth of bone metastasizes. The overall control of bone remodeling is regulated by the binding of RANKL with its receptor or its decoy receptor, respectively, RANK and OPG.



www.acrobiosystems.com