



### Synonym

S1 protein NTD, Spike protein S1 NTD, BetaCoV S1-NTD

### Source

SARS-CoV-2 S1 protein NTD, His Tag (S1D-C52Hc) is expressed from human 293 cells (HEK293). It contains AA Ser 13 - Leu 303 (Accession # [QHD43416.1](#)). The L18F/D80A/D215G/LAL242-244del/R246I mutations were identified on the spike protein N-terminal domain (NTD) in the SARS-CoV-2 Beta variant (Pango lineage: B.1.351; other names: 20H/501Y.V2). Predicted N-terminus: Ser 13

### Molecular Characterization

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 34.5 kDa. The protein migrates as 50-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per µg by the LAL method / rFC method.

### Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:


- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

### SDS-PAGE



SARS-CoV-2 S1 protein NTD, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

### SEC-MALS




The purity of SARS-CoV-2 S1 protein NTD, His Tag (Cat. No. S1D-C52Hc) is more than 90% and the molecular weight of this protein is around 55-65 kDa verified by SEC-MALS.

[Report](#)


### Bioactivity-ELISA

Discounts, Gifts,  
and more!





Immobilized SARS-CoV-2 S1 protein NTD, His Tag (Cat. No. S1D-C52Hc) at 1 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Spike NTD Antibody, Chimeric mAb (Cat. No. SPD-M121) with a linear range of 0.1-2 ng/mL (QC tested).



Immobilized Anti-SARS-CoV-2 Spike NTD Antibody, Chimeric mAb (Cat. No. SPD-M121) at 1 µg/mL (100 µL/well) can bind SARS-CoV-2 S1 protein NTD, His Tag (Cat. No. S1D-C52Hc) with a linear range of 8-63 ng/mL (Routinely tested).

## Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Discounts, Gifts,  
and more!

