

**Human Laminin 511 Protein,
premium grade (Mix & Scale),
(Cat. No. LA1-H5314)
IPSC Suspension Culture Protocol**



Scan this QR code
for special gifts &
more!

1. Experimental instruments and materials

1.1 Instruments: Biosafety cabinet, cell incubator, low temperature horizontal centrifuge, inverted microscope, EnSight (PerkinElmer, HH3400), Microscopic, Flow cytometer.

1.2 Materials: Sterile pipette tips, Sterile EP tube and other consumables.

1.3 Reagents:

Name	Vendor	Cat. No.
mTeSR™-Plus	STEM CELL	100-0276
Y27632	MCE	HY-10071
ReLeSR	STEM CELL	100-0483
PBS	HyClone	SH30256.01
Cell Counting-Lite 2.0 Luminescent Cell Viability Assay	Vazyme	DD1101-02
24 well plate	Nest	702001
OCT4	CST	2750S
NANOG	CST	4903S
SOX2	Santa Cruz	sc-365823
PE-OCT4	Biolegend	653703
PE-SOX2	Biolegend	656103
SSEA4	CST	4755T

1. Experimental instruments and materials


1.4 Activity Assay :

Cell Viability Assay: CellCounting-Lite 2.0 Luminescent Cell Viability Assay

FACS Assay : OCT4, SOX2 , SSEA4

IF Assay : OCT4, NANOG, SOX2

Laminin 511+ ES/iPS Cells



Suspension Method

2. Cell Culture - iPSC

- 2.1. Pre-warm the required volume of cell culture medium containing 10 μ M Y-27632 (final concentration) to 37° C.
- 2.2. Remove the spent culture medium and wash the adherent cells gently with 1 ml of PBS in each well (6 well).
- 2.3. Add 1ml ReLeSR to each well. Stand at room temperature for 6-8 minutes or keep under microscopic observation until cells are not bound to the plate.

NOTE: For the passage of induced pluripotent stem cells (iPSCs), TrypLE Select, ReLeSR, and Accutase are also recommended. When dissociating iPSCs and performing pipetting, the digestion process should be carried out at 37°C for approximately 8 to 10 minutes.
- 2.4. Add equal volume of mTeSR™-Plus, gently mix the cells and transfer it into a centrifuge tube at room temperature at 300g for 5 minutes.
- 2.5. Adjust the concentration of the cell suspension with the pre-warmed medium containing 10 μ M Y-27632.

2.6. Add an adequate volume of laminin 511. The working concentration of 0.55-1 µg/cm² is recommended and should be optimized based on different cell lines.

Culture vessel	Coating concentration	Laminin amount (µg)	mTeSR-plus media
6-well	0.55 µg/cm ²	5.3	2 mL
12-well	0.55 µg/cm ²	2.5	1 mL
24-well	0.55 µg/cm ²	1.1	0.5 mL
35-mm	0.55 µg/cm ²	4.4	2 mL
60-mm	0.55 µg/cm ²	11.6	4 mL
100-mm	0.55 µg/cm ²	30.3	12mL

2.7. Transfer 2 ml of the cell suspension with the laminin 511 to each well. Gently rock the plate back-and-forth and side-to-side to disperse the cells across the surface.

2.8. Incubate the cells for 24 h before exchanging the plating medium for fresh medium (without Y-27632). Replace medium every two days.

3. Cell Counting-Lite 2.0 Luminescent Cell Viability Assay

- 3.1. Pre-warm the 1X Cell Counting-Lite 2.0 Luminescent to room temperature.
- 3.2. Add 250µl 1X Cell Counting-Lite 2.0 Luminescent to 24-well cell culture plate and incubate at room temperature for 15 min.
- 3.3. Transfer 250µl supernatant to a black dark plate and test on the EnSight (PerkinElmer, HH3400).

4. FACS

- 4.1. Collect cells and add 4% paraformaldehyde to fix for 15 min at room temperature.
- 4.2. Add 500µl PBS to wash and incubate at room temperature for 15 min with 350µl 0.5% Triton X-100.
- 4.3. Wash with 500µl PBS after centrifugation and incubate at 4° C with 2% BSA containing antibodies (PE-OCT4, PE-SOX2, SSEA4) for 1h in the dark.
- 4.4. Transfer the cells to a 96-well V-plate with 100µl PBS ready for flow cytometry.

5. IF

- 5.1. Discard the supernatant and add 350µl of 4% paraformaldehyde to fix for 15 min at room temperature (for 24-well plate).
- 5.2. Wash with 500µl PBS and incubate at room temperature with 350µl 0.5% Triton X-100 for 15 min.
- 5.3. Wash with 500µl PBS and incubate at room temperature with 2% BSA for 1h.
- 5.4. Wash with 500µl PBS and incubate overnight at 4°C with 2% BSA coating primary antibody (OCT4, NANOG, SOX2).
- 5.5. Wash 3 times with 500µl PBS and incubate at room temperature with 2% BSA coating second antibody.
- 5.6. Wash 3 times with 500µl PBS and incubate at room temperature with PBS coating DAPI for 10min.
- 5.7. Wash 3 times with 500µl PBS and add 100µl PBS ready for Microscopic observation.

DISCLAIMER: TO THE EXTENT ALLOWED BY LAW, ACROBIOSYSTEMS AND/OR ITS AFFILIATE(S) WILL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, INDIRECT, PUNITIVE, MULTIPLE OR CONSEQUENTIAL DAMAGES IN CONNECTION WITH OR ARISING FROM THIS DOCUMENT, INCLUDING YOUR USE OF IT.

Where and **PROTEINS** **INNOVATION** Advance **BIOMEDICINE**



Cell Therapies



Gene Therapies



Vaccine



Antibody-Drug Conjugates



Therapeutic Antibodies

Discovery & Development

CMC-Manufacturing

CMC- Quality Control

Clinical Studies

Preclinical Studies



Where Proteins and Innovation
Advance Biomedicine



Scan this QR code
for special gifts &
more!